

t66\_tmap\_1  
(TMM46qhDb6NdDQeMbfdsZTavdhf8a2X7qUg)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $m1\_pre\_topc : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_tmap\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_tmap\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc \\
& X0))) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge ((v2\_pre\_topc X1) \wedge (l1\_pre\_topc \\
& X1))) \Rightarrow (\forall X2.((\neg v2\_struct\_0 X2) \wedge (m1\_pre\_topc X2 X1)) \Rightarrow ( \\
& \forall X3.((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 (u1\_struct\_0 X1) \\
& (u1\_struct\_0 X0)) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& (u1\_struct\_0 X1) (u1\_struct\_0 X0)))))) \Rightarrow (\forall X4.((v1\_funct\_1 \\
& X4) \wedge ((v1\_funct\_2 X4 (u1\_struct\_0 X2) (u1\_struct\_0 X0)) \wedge (m1\_subset\_1 \\
& X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (u1\_struct\_0 X2) (u1\_struct\_0 X0)))))) \Rightarrow \\
& ((\forall X5.(m1\_subset\_1 X5 (u1\_struct\_0 X1)) \Rightarrow ((X5 \in u1\_struct\_0 \\
& X2) \Rightarrow (k3\_funct\_2 (u1\_struct\_0 X1) (u1\_struct\_0 X0) X3 X5 = k1\_funct\_1 \\
& X4 X5))) \Rightarrow (r2\_funct\_2 (u1\_struct\_0 X2) (u1\_struct\_0 X0) (k2\_tmap\_1 \\
& X1 X0 X3 X2) X4))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(l1\_pre\_topc X0) \Rightarrow (\forall X1.(m1\_pre\_topc X1 X0) \Rightarrow (l1\_pre\_topc X1)) \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc \\
& X0))) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge ((v2\_pre\_topc X1) \wedge (l1\_pre\_topc \\
& X1)))) \Rightarrow (\forall X2.(m1\_pre\_topc X2 X0) \Rightarrow (\forall X3.(m1\_pre\_topc \\
& X3 X0) \Rightarrow (\forall X4.((v1\_funct\_1 X4) \wedge ((v1\_funct\_2 X4 (u1\_struct\_0 \\
& X2) (u1\_struct\_0 X1)) \wedge (m1\_subset\_1 X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& (u1\_struct\_0 X2) (u1\_struct\_0 X1)))))) \Rightarrow ((m1\_pre\_topc X3 X2) \Rightarrow \\
& (k3\_tmap\_1 X0 X1 X2 X3 X4 = k2\_partfun1 (u1\_struct\_0 X2) (u1\_struct\_0 \\
& X1) X4 (u1\_struct\_0 X3))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc \\
& X0))) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge ((v2\_pre\_topc X1) \wedge (l1\_pre\_topc \\
& X1)))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 (u1\_struct\_0 \\
& X0) (u1\_struct\_0 X1)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& (u1\_struct\_0 X0) (u1\_struct\_0 X1)))))) \Rightarrow (\forall X3.(m1\_pre\_topc \\
& X3 X0) \Rightarrow (k2\_tmap\_1 X0 X1 X2 X3 = k2\_partfun1 (u1\_struct\_0 X0) (u1\_struct\_0 \\
& X1) X2 (u1\_struct\_0 X3))))
\end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v2\_pre\_topc X0) \wedge (l1\_pre\_topc X0)) \Rightarrow (\forall X1. \\
& (m1\_pre\_topc X1 X0) \Rightarrow (v2\_pre\_topc X1))
\end{aligned} \tag{5}$$

**Theorem 1**

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc \\
& X0))) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge ((v2\_pre\_topc X1) \wedge (l1\_pre\_topc \\
& X1)))) \Rightarrow (\forall X2.((\neg v2\_struct\_0 X2) \wedge (m1\_pre\_topc X2 X0)) \Rightarrow ( \\
& \forall X3.((\neg v2\_struct\_0 X3) \wedge (m1\_pre\_topc X3 X0)) \Rightarrow (\forall X4. \\
& ((v1\_funct\_1 X4) \wedge ((v1\_funct\_2 X4 (u1\_struct\_0 X3) (u1\_struct\_0 \\
& X1)) \wedge (m1\_subset\_1 X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (u1\_struct\_0 \\
& X3) (u1\_struct\_0 X1)))))) \Rightarrow ((m1\_pre\_topc X2 X3) \Rightarrow (\forall X5.( \\
& (v1\_funct\_1 X5) \wedge ((v1\_funct\_2 X5 (u1\_struct\_0 X2) (u1\_struct\_0 \\
& X1)) \wedge (m1\_subset\_1 X5 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (u1\_struct\_0 \\
& X2) (u1\_struct\_0 X1)))))) \Rightarrow ((\forall X6.(m1\_subset\_1 X6 (u1\_struct\_0 \\
& X3)) \Rightarrow ((X6 \in u1\_struct\_0 X2) \Rightarrow (k3\_funct\_2 (u1\_struct\_0 X3) (u1\_struct\_0 \\
& X1) X4 X6 = k1\_funct\_1 X5 X6))) \Rightarrow (r2\_funct\_2 (u1\_struct\_0 X2) (u1\_struct\_0 \\
& X1) (k3\_tmap\_1 X0 X1 X3 X2 X4) X5))))))
\end{aligned}$$