

t66\_xxreal\_1

(TMNKws7gUMNnqAgLFme7HTUjE6rXYpogB61)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (\forall X3.(v1\_xxreal\_0 X3) \Rightarrow (((r1\_xxreal\_0 \\ & X0 X1) \wedge (r1\_tarski (k1\_xxreal\_1 X0 X1) (k1\_xxreal\_1 X2 X3))) \Rightarrow (( \\ & r1\_xxreal\_0 X2 X0) \wedge (r1\_xxreal\_0 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\ & (r1\_xxreal\_0 X0 X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ & (r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X0)) \Rightarrow (X0 = X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.r1\_tarski X0 X0 \quad (4)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (\forall X3.(v1\_xxreal\_0 X3) \Rightarrow (((r1\_xxreal\_0 \\ & X0 X1) \wedge (k1\_xxreal\_1 X0 X1 = k1\_xxreal\_1 X2 X3)) \Rightarrow ((X0 = X2) \wedge (X1 = X3)))))) \end{aligned}$$