

t67_complfld
(TMPkh1fus9b7MmDbhMy3WdwKZY6mhbDboGU)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_complfld : \iota$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k5_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow ((k17_complex1 (k6_xcmplx_0 X0 X1) = k6_numbers) \Leftrightarrow (X0 = X1))) \quad (1)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 k1_complfld)) \Rightarrow (\forall X2.(v1_xcmplx_0 X2) \Rightarrow (\forall X3.(v1_xcmplx_0 X3) \Rightarrow (((X0 = X2) \wedge (X1 = X3)) \Rightarrow (k5_algstr_0 k1_complfld X0 X1 = k4_binop_2 X2 X3)))))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k4_binop_2 X0 X1 = k6_xcmplx_0 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (v1_xcmplx_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 k1_complfld)) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 k1_complfld)) \Rightarrow ((k17_complex1 (k5_algstr_0 k1_complfld X0 X1) = k6_numbers) \Leftrightarrow (X0 = X1)))$$