

# t67\_finseq\_2 (TMJQKWGQeQttTYBR- FrhZYx1nwrsUN6McW3z)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((r1\_tarski (k2\_zfmisc\_1 (k1\_tarski X0) (k10\_xtuple\_0 X1)) (k9\_xtuple\_0 X2)) \Rightarrow (k9\_xtuple\_0 (k5\_funcop\_1 X2 X0 X1) = k4\_finseq\_1 X1))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (3)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1. (m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow ((X1 = k3\_finseq\_1 X0) \Leftrightarrow (k2\_finseq\_1 X1 = k9\_xtuple\_0 X0))) \quad (4)$$

## Theorem 1

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2))) \Rightarrow (\forall X3. ((v1\_relat\_1 X3) \wedge (v1\_funct\_1 X3)) \Rightarrow (((r1\_tarski (k2\_zfmisc\_1 (k1\_tarski X0) (k10\_xtuple\_0 X1)) (k9\_xtuple\_0 X3)) \wedge (X2 = k5\_funcop\_1 X3 X0 X1)) \Rightarrow (k3\_finseq\_1 X2 = k3\_finseq\_1 X1))))$$