

## t67\_finseq\_6

(TMHjLDH6BtD7jNzsj1kGcTSBehhWQL3oMdw)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_finseq\_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X2)) \wedge (v1\_xboole\_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 X0) \Rightarrow \\ & (\forall X2. (m2\_finseq\_1 X2 X0) \Rightarrow ((X1 \in k10\_xtuple\_0 X2) \Rightarrow (k1\_finseq\_5 \\ & X0 X2 X1 = k7\_finseq\_1 (k5\_finseq\_4 X2 X1) (k12\_finseq\_1 X0 X1)))))) \quad (3) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ & X1)))) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 \\ & X2)))) \Rightarrow (k7\_finseq\_1 (k7\_finseq\_1 X0 X1) X2 = k7\_finseq\_1 X0 (k7\_finseq\_1 \\ & X1 X2)))) \quad (4) \end{aligned}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ X1))) \Rightarrow (k10\_xtuple\_0 (k7\_finseq\_1 X0 X1) = k2\_xboole\_0 (k10\_xtuple\_0 \\ X0) (k10\_xtuple\_0 X1))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ X1))) \Rightarrow (\forall X2. (X2 \in k6\_subset\_1 (k10\_xtuple\_0 X0) (k10\_xtuple\_0 \\ X1)) \Rightarrow (k5\_finseq\_4 (k7\_finseq\_1 X1 X0) X2 = k7\_finseq\_1 X1 (k5\_finseq\_4 \\ X0 X2)))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_finseq\_1 X1 X0) \wedge (m1\_finseq\_1 X2 X0)) \Rightarrow (k8\_finseq\_1 X0 X1 X2 = k7\_finseq\_1 X1 X2) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X1 X0)) \Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))) \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \quad (12)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_finseq\_1 X1 X0) \wedge (m1\_finseq\_1 X2 X0)) \Rightarrow (m2\_finseq\_1 (k8\_finseq\_1 X0 X1 X2) X0) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.m1\_subset\_1 (k6\_subset\_1 X0 X1) (k1\_zfmisc\_1 X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow((v1\_relat\_1 (k5\_finseq\_4 X0 X1))\wedge((v1\_funct\_1 (k5\_finseq\_4 X0 X1))\wedge(v1\_finseq\_1 (k5\_finseq\_4 X0 X1)))) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(m1\_finseq\_1 X1 X0))\Rightarrow(m2\_finseq\_1 (k1\_finseq\_5 X0 X1 X2) X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow(m2\_finseq\_1 (k12\_finseq\_1 X0 X1) X0) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k2\_xboole\_0 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in X0)\vee(X3 \in X1))) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.k2\_xboole\_0 X0 X1 = k2\_xboole\_0 X1 X0 \quad (19)$$

**Theorem 1**

$$\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow(\forall X2.(m2\_finseq\_1 X2 X0)\Rightarrow(\forall X3.(m2\_finseq\_1 X3 X0)\Rightarrow((X1 \in k6\_subset\_1 (k10\_xtuple\_0 X2) (k10\_xtuple\_0 X3))\Rightarrow(k1\_finseq\_5 X0 (k8\_finseq\_1 X0 X3 X2) X1 = k8\_finseq\_1 X0 X3 (k1\_finseq\_5 X0 X2 X1))))))$$