

t67_xreal_1 (TM-
MxbzvUX3SwXE9fXQZB7QqhJ8QdRxB9ZzE)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X2 k6_numbers)) \Rightarrow \\ & (r1_xxreal_0 (k3_xcmplx_0 X1 X2) (k3_xcmplx_0 X0 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg \\ & r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 X1 (k6_xcmplx_0 X1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 \\ & X1 X2)) \Rightarrow (r1_xxreal_0 (k2_xcmplx_0 X1 X0) X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 (k6_xcmplx_0 X1 X2)) \Rightarrow (r1_xxreal_0 \\ & X2 (k6_xcmplx_0 X1 X0)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow (\\ & r1_xxreal_0 X0 X2)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k2_xcmplx_0 X0 X1) X2) \Rightarrow (r1_xxreal_0 \\ & X0 (k6_xcmplx_0 X2 X1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k3_xcmplx_0 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0)\wedge(v1_xcmplx_0 X1))\Rightarrow(k3_xcmplx_0 X0 X1 = k3_xcmplx_0 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(\forall X2. \\ & (v1_xreal_0 X2)\Rightarrow(\forall X3.(v1_xreal_0 X3)\Rightarrow(((r1_xxreal_0 \\ & X0 k6_numbers)\wedge((r1_xxreal_0 X1 X0)\wedge((r1_xxreal_0 X2 k6_numbers)\wedge \\ & (r1_xxreal_0 X3 X2))))\Rightarrow(r1_xxreal_0 (k3_xcmplx_0 X0 X2) (k3_xcmplx_0 \\ & X1 X3)))))) \end{aligned}$$