

t68\_afinsq\_1 (TMR-  
Jfdm6BMdvtqcdY1MYEPNuz5FpLTb11kz)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_afinsq\_1 : \iota \Rightarrow o$  be given. Let  $k2\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_card\_1 : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k1\_card\_1 : \iota \Rightarrow \iota$  be given. Let  $v5\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_card\_1 : \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v7\_ordinal1 X0) \Rightarrow (X0 = ReplSep (toset (\lambda X1 : \iota. \\ m2\_subset\_1 X1 k1\_numbers k5\_numbers)) (\lambda X1 : \iota. \neg r1\_xxreal\_0 \\ X0 X1) (\lambda X1 : \iota. X1)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (\forall X2. (X2 \in X0) \Leftrightarrow (X2 \in X1)) \Rightarrow (X0 = X1) \quad (2)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.(v1\_finset\_1 X0) \Rightarrow (k5\_card\_1 X0 = k1\_card\_1 X0) \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 \\ X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (k2\_afinsq\_1 X0 = k9\_xtuple\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 \\ X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (k1\_card\_1 X0 = k9\_xtuple\_0 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_finset\_1 X0) \Rightarrow ((v1\_finset\_1 (k1\_card\_1 X0)) \wedge (v1\_card\_1 (k1\_card\_1 X0))) \quad (7)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge (v1\_afinsq\_1 X0))))) \Rightarrow ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge (v1\_funct\_1 X0))) \quad (8)$$

Assume the following.

$$\forall X0.((v3\_ordinal1 X0) \wedge (v1\_finset\_1 X0)) \Rightarrow (v7\_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1\_card\_1 X0) \Rightarrow (v3\_ordinal1 X0) \quad (10)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge (v1\_afinsq\_1 X0))))) \Rightarrow (k2\_afinsq\_1 X0 = \text{ReplSep } (\text{toset } (\lambda X1 : \iota.m2\_subset\_1 X1 k1\_numbers k5\_numbers)) (\lambda X1 : \iota.\neg r1\_xxreal\_0 (k5\_card\_1 X0) X1) (\lambda X1 : \iota.X1)))$$