

t68_finseq_4 (TMFBGdFK- WynsQ7tSp2HnRWiy3SCkVuWfJKf)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow (\forall X2.(X2 \in k1_relset_1 k5_numbers X0) \Rightarrow (X2 \in k1_relset_1 \\ & k5_numbers (k7_finseq_1 X0 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_finseq_1 X1 X0) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k8_finseq_1 X0 X1 X2) X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))))\Rightarrow(((v1_relat_1 (k7_finseq_1 X0 X1))\wedge((v1_funct_1 (k7_finseq_1 X0 X1))\wedge(v1_finseq_1 (k7_finseq_1 X0 X1)))))) \quad (9)$$

Assume the following.

$$\forall X0.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow((\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow(\forall X2.((v1_relat_1 X2)\wedge((v1_funct_1 X2)\wedge(v1_finseq_1 X2)))\Rightarrow((X2 = k7_finseq_1 X0 X1)\Leftrightarrow((k4_finseq_1 X2 = k2_finseq_1 (k2_nat_1 (k3_finseq_1 X0) (k3_finseq_1 X1))))\wedge((\forall X3.(v7_ordinal1 X3)\Rightarrow((X3 \in k4_finseq_1 X0)\Rightarrow(k1_funct_1 X2 X3 = k1_funct_1 X0 X3))\wedge(\forall X3.(v7_ordinal1 X3)\Rightarrow((X3 \in k4_finseq_1 X1)\Rightarrow(k1_funct_1 X2 (k2_nat_1 (k3_finseq_1 X0) X3) = k1_funct_1 X1 X3)))))))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (13)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.(m2_finseq_1 X2 X1)\Rightarrow(\forall X3.(m2_finseq_1 X3 X1)\Rightarrow((X0 \in k4_finseq_1 X2)\Rightarrow(k7_partfun1 X1 (k8_finseq_1 X1 X2 X3) X0 = k7_partfun1 X1 X2 X0))))))$$