

t68_pnproc_1
(TMWY1k7MUVn8pkxkPCCMY7wMHn8EYVwTSrE)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_pnproc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 (k2_xboole_0 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow (k7_finseq_1 X0 X1 = k2_xboole_0 X0 (k12_pnproc_1 X1 (k3_finseq_1 \\ & X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow (r1_tarski (k12_pnproc_1 X1 (k3_finseq_1 X0)) (k7_finseq_1 \\ & X0 X1))) \end{aligned}$$