

t68_qc_lang2
(TMGgDgxJrqizr9efjkCimpbhesaLSpQdFJv)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $r3_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_qc_lang2 : \iota \Rightarrow \iota$ be given. Let $r2_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k13_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0)) \Rightarrow ((r2_qc_lang2 \\ X0 X1 X2) \Leftrightarrow (r3_qc_lang2 X0 X1 (k13_qc_lang1 X0 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k12_qc_lang1 X0) (k9_qc_lang1 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (k1_qc_lang2 X0 = k13_qc_lang1 X0 (k12_qc_lang1 X0)) \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((r3_qc_lang2 X0 X1 (k1_qc_lang2 X0)) \Leftrightarrow (r2_qc_lang2 X0 X1 \\ (k12_qc_lang1 X0)))) \end{aligned}$$