

t69_arytm_3
(TMT4DFx9YuBeT2oPvabv1CQtR2RTGXpE5P2)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_arytm_3 : \iota$ be given. Let $r3_arytm_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_arytm_3) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k5_arytm_3) \Rightarrow (\forall X2.(m1_subset_1 X2 k5_arytm_3) \Rightarrow (((r3_arytm_3 \\ & X0 X1) \wedge (r3_arytm_3 X1 X2)) \Rightarrow (r3_arytm_3 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_subset_1 X0 k5_arytm_3) \wedge (m1_subset_1 \\ & X1 k5_arytm_3)) \Rightarrow ((r3_arytm_3 X0 X1) \vee (r3_arytm_3 X1 X0)) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_arytm_3) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k5_arytm_3) \Rightarrow (\forall X2.(m1_subset_1 X2 k5_arytm_3) \Rightarrow (\neg((\\ & (\neg r3_arytm_3 X1 X0) \wedge (r3_arytm_3 X1 X2)) \vee ((r3_arytm_3 X0 X1) \wedge (\\ & \neg r3_arytm_3 X2 X1))) \wedge (r3_arytm_3 X2 X0)))) \end{aligned}$$