

t69_complex1
(TMZWFHy4SVnGdULPEASDttocAEnXkXdktpN)

October 27, 2020

Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k15_complex1 : \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k14_complex1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k17_complex1 (k3_xcmplx_0 X0 X1) = k8_real_1 (k17_complex1 X0) (k17_complex1 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k17_complex1 (k15_complex1 X0) = k17_complex1 X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k15_complex1 X0 = k14_complex1 X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (v1_xcmplx_0 (k14_complex1 X0)) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k17_complex1 (k3_xcmplx_0 X0 X0) = k17_complex1 (k3_xcmplx_0 X0 (k15_complex1 X0)))$$