

t69_compos_1
(TMWJgBMwYo9NZxSD6Nz1HrHimt4VQyu1otG)

October 27, 2020

Let $l1_compos_1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_compos_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_afinsq_1 : \iota \Rightarrow o$ be given. Let $k63_valued_1 : \iota \Rightarrow \iota$ be given. Let $k10_compos_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((l1_compos_1 X0) \wedge ((v1_relat_1 X1) \wedge ((\\ & v4_relat_1 X1 k5_numbers) \wedge ((v5_relat_1 X1 (u1_compos_1 X0)) \wedge \\ & ((v1_funct_1 X1) \wedge ((v1_finset_1 X1) \wedge (v1_afinsq_1 X1)))))) \Rightarrow \\ & (k63_valued_1 (k10_compos_1 X0 X1) = X1) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0. (l1_compos_1 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 k5_numbers) \wedge ((v5_relat_1 X1 \\ & (u1_compos_1 X0)) \wedge ((v1_funct_1 X1) \wedge ((v1_finset_1 X1) \wedge (v1_afinsq_1 \\ & X1)))))) \Rightarrow (k63_valued_1 (k10_compos_1 X0 X1) = X1)) \end{aligned}$$