

# t69\_flang\_3 (TMaSHFfNkmnAcy- DtMx8K7S1QGEQZvGusJ7d)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_flang\_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (k2\_flang\_3 X0 X1 = k1\_flang\_3 X0 X1 np\_1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k8\_afinsq\_1 X1))) \Rightarrow (\forall X3. (v7\_ordinal1 X3) \Rightarrow ((k1\_flang\_3 X1 X2 X3 = k1\_tarski X0) \Rightarrow (X0 = k2\_flang\_1 X1))) \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. (m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (5)$$

**Theorem 1**

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k8\_afinsq\_1 X1))) \Rightarrow ((k2\_flang\_3 X1 X2 = k1\_tarski X0) \Rightarrow (X0 = k2\_flang\_1 X1))$$