

t6_bvfunc26 (TMartDfP- bYAYX93jDwwsPEmLKd2fH18WKRg)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_bvfunc26 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ & (k5_bvfunc26 X0 X1 X2) (k1_bvfunc_1 X0 (k2_bvfunc_1 X0 X1 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))))) \wedge ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 \\ & X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow ((r2_funct_2 X0 X1 X2 \\ & X3) \Leftrightarrow (X2 = X3)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (k1_bvfunc_1 X0 (k1_bvfunc_1 \\ & X0 X1) = X1) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow (k2_bvfunc_1 X0 X1 X1 = X1) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k5_bvfunc26 X0 X1 X2)) \wedge (\\ & (v1_funct_2 (k5_bvfunc26 X0 X1 X2) X0 k6_margrel1) \wedge (m1_subset_1 \\ & (k5_bvfunc26 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k1_bvfunc_1 \\ & X0 X1)) \wedge ((v1_funct_2 (k1_bvfunc_1 X0 X1) X0 k6_margrel1) \wedge (m1_subset_1 \\ & (k1_bvfunc_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((r2_funct_2 X0 k6_margrel1 \\ & (k5_bvfunc26 X0 X1 X1) (k1_bvfunc_1 X0 X1)) \wedge (r2_funct_2 X0 k6_margrel1 \\ & (k1_bvfunc_1 X0 (k5_bvfunc26 X0 X1 X1)) X1)) \end{aligned}$$