

t6_comptrig
(TMF7PyPTVB65qG62EqbnbKoYVRrRa2DXTRK)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ & (r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & k2_rcomp_1 X0 X1 = k4_xxreal_1 X0 X1) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (\neg (X3 \in k2_rcomp_1 \\ & X0 X2) \wedge (\neg X3 \in k2_rcomp_1 X0 X1) \wedge ((X3 \neq X1) \wedge (\neg X3 \in k2_rcomp_1 X1 X2))))))) \end{aligned}$$