

t6_comseq_1
(TMWfy1C6wDNNpZVSDhQoN9BRwku5yJXB9zo)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(X0 \neq k1_xboole_0) \Rightarrow (\forall X1.\exists X2.((v1_relat_1 \\ X2) \wedge (v1_funct_1 X2)) \wedge ((k9_xtuple_0 X2 = X0) \wedge (k10_xtuple_0 X2 = \\ k1_tarski X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(v1_relat_1 X2) \Rightarrow (((r1_tarski \\ (k9_xtuple_0 X2) X0) \wedge (r1_tarski (k10_xtuple_0 X2) X1)) \Rightarrow (m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.r1_tarski X0 X0 \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (\\ k2_relset_1 X0 X1 = k10_xtuple_0 X1) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(\\ k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k2_numbers \quad (9)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow\\ (X2 \in X1)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow\\ (X2 = X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1\\ (k2_zfmisc_1 X0 X1)))\Rightarrow(((X1\neq k1_xboole_0)\Rightarrow((v1_funct_2 X2 X0\\ X1)\Leftrightarrow(X0 = k1_relset_1 X0 X2)))\wedge((X1 = k1_xboole_0)\Rightarrow((v1_funct_2\\ X2 X0 X1)\Leftrightarrow(X2 = k1_xboole_0)))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1\\ (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1\\ (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (15)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k2_numbers)\Rightarrow(\exists X1.((v1_funct_1\\ X1)\wedge((v1_funct_2 X1 k5_numbers k2_numbers)\wedge(m1_subset_1 X1 (\\ k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers))))\wedge(k2_relset_1\\ k2_numbers X1 = k1_tarski X0))$$