

t6_cqc_lang
(TMV5ZTP1mg6S1Y4Qwf4t3aQhL17HK1cVyNM)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $k6_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k5_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (k6_qc_lang3 X0 (k12_qc_lang1 X0) = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (k5_qc_lang3 X0 (k12_qc_lang1 X0) = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 X0)) \Rightarrow ((m2_subset_1 X1 (k9_qc_lang1 X0) (k3_cqc_lang X0)) \Leftrightarrow ((k6_qc_lang3 X0 X1 = k1_xboole_0) \wedge (k5_qc_lang3 X0 X1 = k1_xboole_0)))) \quad (3)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k12_qc_lang1 X0) (k9_qc_lang1 X0)) \quad (4)$$

Theorem 1

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m2_subset_1 (k12_qc_lang1 X0) (k9_qc_lang1 X0) (k3_cqc_lang X0))$$