

t6_fcont_1

(TMX8eNHpTrgx1uD6WN3YjHogPY4bfyXWQo8)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_4 : \iota \Rightarrow \iota$ be given. Let $r1_fcont_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers)))) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_fcont_1 \\ X0 X1) \Leftrightarrow (\forall X2.(m1_rcomp_1 X2 (k1_seq_1 X0 X1)) \Rightarrow (\exists X3. \\ (m1_rcomp_1 X3 X1) \wedge (r1_tarski (k7_relset_1 k1_numbers k1_numbers \\ X0 X3) X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X0 \in \\ k9_xtuple_0 X1) \Rightarrow (k9_relat_1 X1 X0 = k1_tarski (k1_funct_1 X1 X0))) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski \\ X0 X1) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \tag{4}$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\forall X1. (m1_rcomp_1 X1 X0) \Rightarrow (X0 \in X1)) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1)\Rightarrow(k7_relat_1 X1 X0 = k7_relat_1 X1 (k3_xboole_0 (k9_xtuple_0 X1) X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.r1_tarski X0 X0 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0))\Rightarrow(k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(k1_seq_4 X0 = k1_tarski X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(v1_xreal_0 (k1_funct_1 X0 X1)) \quad (13)$$

Assume the following.

$$v3_membered k1_numbers \quad (14)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(m1_rcomp_1 X1 X0)\Rightarrow(m1_subset_1 X1 (k1_zfmisc_1 k1_numbers))) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0))\Rightarrow(m1_subset_1 (k9_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \quad (16)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.k9_relat_1 X0 X1 = k7_relat_1 X0 (k1_tarski X1)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v3_valued_0 X2)) \quad (21)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow ((\exists X2.(m1_rcomp_1 X2 X0) \wedge (k9_subset_1 k1_numbers (k1_relset_1 k1_numbers X1) X2 = k1_seq_4 X0)) \Rightarrow (r1_fcont_1 X1 X0)))$$