

t6_finseq_5

(TMUxnbt4HkCVeS2NF8cSwGc4M3rJS5QzMLM)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_card_1 : \iota \Rightarrow o$ be given. Let $k16_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_tarski (k2_finseq_1 X0) (k2_finseq_1 X1)))) \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((X0 \neq k1_xboole_0) \Leftrightarrow (r1_xxreal_0 np_1 (k3_finseq_1 X0))) \quad (4)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.r1_tarski\ X0\ X0 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow (k4_finseq_1\ X0 = k9_xtuple_0\ X0) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow (k3_finseq_1\ X0 = k1_card_1\ X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1\ X1)\wedge(v4_relat_1\ X1\ X0))\Rightarrow (k1_relset_1\ X0\ X1 = k9_xtuple_0\ X1) \quad (10)$$

Assume the following.

$$\forall X0.(v1_finset_1\ X0)\Rightarrow((v1_finset_1\ (k1_card_1\ X0))\wedge(v1_card_1\ (k1_card_1\ X0))) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow ((v1_relat_1\ (k16_finseq_1\ k6_numbers\ X0))\wedge((v1_funct_1\ (k16_finseq_1\ k6_numbers\ X0))\wedge((v1_xboole_0\ (k16_finseq_1\ k6_numbers\ X0))\wedge(v1_finseq_1\ (k16_finseq_1\ k6_numbers\ X0)))))) \quad (12)$$

Assume the following.

$$\forall X0.v1_card_1\ (k1_card_1\ X0) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (14)$$

Assume the following.

$$\forall X0.((v3_ordinal1\ X0)\wedge(v1_finset_1\ X0))\Rightarrow(v7_ordinal1\ X0) \quad (15)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow ((v1_relat_1\ X0)\wedge((v4_relat_1\ X0\ k5_numbers)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))) \quad (16)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0))) \quad (17)$$

Assume the following.

$$\forall X0.(v1_card_1 X0) \Rightarrow (v3_ordinal1 X0) \quad (18)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0) \wedge ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow ((np_1 \in k4_finseq_1 X0) \wedge (k3_finseq_1 X0 \in k4_finseq_1 X0))$$