

t6_fscirc_1

(TMGTV68gyUyTQprH2AMBA8nrnkb6vDAkrnu)

October 27, 2020

Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k4_fscirc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k11_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v11_struct_0 : \iota \Rightarrow o$ be given. Let $v1_msualg_1 : \iota \Rightarrow o$ be given. Let $k3_fscirc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_circcomb : \iota \Rightarrow o$ be given. Let $v2_circcomb : \iota \Rightarrow o$ be given. Let $v3_circcomb : \iota \Rightarrow o$ be given. Let $k4_facirc_1 : \iota$ be given. Let $k3_twoscomp : \iota$ be given. Let $k2_twoscomp : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (X0 \in u1_struct_0 (k5_circcomb \\ & X2 (k10_finseq_1 X0 X1))) \wedge ((X1 \in u1_struct_0 (k5_circcomb X2 (k10_finseq_1 \\ & X0 X1))) \wedge (k4_tarski (k10_finseq_1 X0 X1) X2 \in u1_struct_0 (k5_circcomb \\ & X2 (k10_finseq_1 X0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\ & ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow (\forall X2. (m1_subset_1 \\ & X2 (u1_struct_0 X0)) \Rightarrow ((X2 \in u1_struct_0 (k2_circcomb X0 X1)) \wedge (\\ & X2 \in u1_struct_0 (k2_circcomb X1 X0)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 (k11_finseq_1 X0 X1 X2)) \wedge (v1_funct_1 (k11_finseq_1 X0 X1 X2)) \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow ((\neg v2_struct_0 (k5_circcomb X0 X1)) \wedge ((\neg v11_struct_0 (\\ & k5_circcomb X0 X1)) \wedge (v1_msualg_1 (k5_circcomb X0 X1)))) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 (k10_finseq_1 X0 X1))\wedge(v1_funct_1 (k10_finseq_1 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.v1_finseq_1 (k11_finseq_1 X0 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.v1_finseq_1 (k10_finseq_1 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow((\neg v11_struct_0 (k5_circcomb X0 X1))\wedge((v1_msualg_1 (k5_circcomb X0 X1))\wedge(l1_msualg_1 (k5_circcomb X0 X1)))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(\neg v2_struct_0 (k3_fscirc_1 X0 X1 X2))\wedge((\neg v11_struct_0 (k3_fscirc_1 X0 X1 X2))\wedge((v1_msualg_1 (k3_fscirc_1 X0 X1 X2))\wedge((v1_circcomb (k3_fscirc_1 X0 X1 X2))\wedge((v2_circcomb (k3_fscirc_1 X0 X1 X2))\wedge((v3_circcomb (k3_fscirc_1 X0 X1 X2))\wedge(l1_msualg_1 (k3_fscirc_1 X0 X1 X2)))))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v2_struct_0 X0)\wedge(l1_msualg_1 X0))\wedge((\neg v2_struct_0 X1)\wedge(l1_msualg_1 X1)))\Rightarrow((\neg v2_struct_0 (k2_circcomb X0 X1))\wedge((v1_msualg_1 (k2_circcomb X0 X1))\wedge(l1_msualg_1 (k2_circcomb X0 X1)))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k4_fscirc_1 X0 X1 X2 = k2_circcomb (k3_fscirc_1 X0 X1 X2) (k5_circcomb k4_facirc_1 (k11_finseq_1 (k4_tarski (k10_finseq_1 X0 X1) k3_twoscomp) (k4_tarski (k10_finseq_1 X1 X2) k2_twoscomp) (k4_tarski (k10_finseq_1 X0 X2) k3_twoscomp))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k3_fscirc_1 X0 X1 X2 = k2_circcomb (k2_circcomb (k5_circcomb k3_twoscomp (k10_finseq_1 X0 X1)) (k5_circcomb k2_twoscomp (k10_finseq_1 X1 X2))) (k5_circcomb k3_twoscomp (k10_finseq_1 X0 X2)) \quad (13)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.(X0 \in u1_struct_0 (k4_fscirc_1 X0 X1 X2)) \wedge ((X1 \in u1_struct_0 (k4_fscirc_1 X0 X1 X2)) \wedge (X2 \in u1_struct_0 (k4_fscirc_1 X0 X1 X2)))$$