

t6_grfunc_1
(TMF_{x6jBviyg75udECcW97JJs9Cqqf8Yj3Ue})

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((k4_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (X1 = k1_funct_1 X2 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X2 = k1_tarski (k4_tarski X0 X1)) \Rightarrow (k1_funct_1 X2 X0 = X1))$$