

t6_integra6 (TM-
Fus5BCTN9wdRZqEUMGosHdnjhaWn1Z9zF)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_integral : \iota \Rightarrow \iota$ be given. Let $k3_integral5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xxreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k3_integral (k3_integral5 X0 X1) = k6_xcmplx_0 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 k6_numbers (k6_xcmplx_0 X1 X0)))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (r1_xxreal_0 X0 (k4_xxreal_0 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (k18_complex1 (k6_xcmplx_0 X0 X1) = k18_complex1 (k6_xcmplx_0 X1 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k18_complex1 X0 = k16_complex1 X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k6_xcmplx_0 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k3_xxreal_0 X0 X1 = X0)) \wedge ((\neg r1_xxreal_0 X0 X1) \Rightarrow (k3_xxreal_0 X0 X1 = X1)))) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = X0)) \wedge ((\neg r1_xxreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = k4_xcmplx_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X1 X0) \Rightarrow (k4_xxreal_0 X0 X1 = X0)) \wedge ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k4_xxreal_0 X0 X1 = X1)))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k4_xxreal_0 X0 X1 = k4_xxreal_0 X1 X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k3_xxreal_0 X0 X1 = k3_xxreal_0 X1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (13)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (k3_integra1 (k3_integra5 (k3_xxreal_0 X0 X1) (k4_xxreal_0 X0 X1)) = k18_complex1 (k6_xcmplx_0 X1 X0))))$$