

t6_kurato_0 (TMNDc-
cXwGeXz7mhfMVfUJYm233ByYkJUmbM)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_kurato_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_kurato_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (\forall X2. (X2 \in k4_kurato_0 \\ & X0 X1) \Leftrightarrow (\forall X3. (m1_subset_1 X3 k5_numbers) \Rightarrow (\exists X4. (\\ & m1_subset_1 X4 k5_numbers) \wedge (X2 \in k3_funct_2 k5_numbers (k9_setfam_1 \\ & X0) X1 (k2_nat_1 X3 X4)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (\forall X2. (X2 \in k3_kurato_0 \\ & X0 X1) \Leftrightarrow (\exists X3. (m1_subset_1 X3 k5_numbers) \wedge (\forall X4. (\\ & m1_subset_1 X4 k5_numbers) \Rightarrow (X2 \in k3_funct_2 k5_numbers (k9_setfam_1 \\ & X0) X1 (k2_nat_1 X3 X4)))))) \end{aligned} \tag{2}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_nat_1 X1 X0) \tag{5}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (6)$$

Theorem 1

$$\forall X0.\forall X1.((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (r1_tarski (k3_kurato_0 X0 X1) (k4_kurato_0 X0 X1))$$