

# t6\_mboolean (TMXFB- HCVX1RSKpvBC99MUUcRcnTrppfC8RW)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r2\_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_mboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v2\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Rightarrow (r1\_tarski (k1\_zfmisc\_1 X0) (k1\_zfmisc\_1 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow ((v1\_relat\_1 (k1\_mboolean X0 X1)) \wedge ((v2\_relat\_1 (k1\_mboolean X0 X1)) \wedge ((v4\_relat\_1 (k1\_mboolean X0 X1) X0) \wedge ((v1\_funct\_1 (k1\_mboolean X0 X1)) \wedge (v1\_partfun1 (k1\_mboolean X0 X1) X0)))))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow ((r2\_pboole X0 X1 X2) \Leftrightarrow (\forall X3. (X3 \in X0) \Rightarrow (r1\_tarski (k1\_funct\_1 X1 X3) (k1\_funct\_1 X2 X3)))))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow ((X2 = k1\_mboolean X0 X1) \Leftrightarrow (\forall X3. (X3 \in X0) \Rightarrow (k1\_funct\_1 X2 X3 = k1\_zfmisc\_1 (k1\_funct\_1 X1 X3)))))) \quad (4)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow \\ & ((r2\_pboole X0 X1 X2) \Rightarrow (r2\_pboole X0 (k1\_mboolean X0 X1) (k1\_mboolean \\ & X0 X2)))) \end{aligned}$$