

t6_nat_2

(TMZ597nCNHmzszTZxQ4v5S1imBz14mqrLGC)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\forall X1. (v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k1_xreal_0 X1 X0 = k6_xcmplx_0 X1 X0))) \quad (4)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X0 \in k2_finseq_1 X1) \Leftrightarrow ((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 X1)))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k2_xcmplx_0 X0 X1) X2) \Leftrightarrow (r1_xxreal_0 \\ & X0 (k6_xcmplx_0 X2 X1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k6_xcmplx_0 X0 X3) (\\ & k6_xcmplx_0 X1 X2)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (r1_xxreal_0 X0 (k2_xcmplx_0 X0 X1))) \quad (8)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (k7_nat_d X0 X1 = k1_xreal_0 X0 X1) \quad (10)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k6_xcmplx_0 X0 X1)) \quad (13)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (14)$$

Assume the following.

$$v3_membered k1_numbers \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(v7_ordinal1\ X1))\Rightarrow(v7_ordinal1\ (k2_xcmplx_0\ X0\ X1)) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(v7_ordinal1\ X1))\Rightarrow(m1_subset_1\ (k7_nat_d\ X0\ X1)\ k5_numbers) \quad (17)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (18)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(m1_subset_1\ (k2_finseq_1\ X0)\ (k1_zfmisc_1\ k5_numbers)) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0\ X0)\wedge(v1_xxreal_0\ X1))\Rightarrow((r1_xxreal_0\ X0\ X1)\vee(r1_xxreal_0\ X1\ X0)) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0\ X0)\wedge(v1_xcmplx_0\ X1))\Rightarrow(k2_xcmplx_0\ X0\ X1 = k2_xcmplx_0\ X1\ X0) \quad (21)$$

Assume the following.

$$\forall X0.(v3_membered\ X0)\Rightarrow(v1_membered\ X0) \quad (22)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xxreal_0\ X0) \quad (23)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xreal_0\ X0) \quad (24)$$

Assume the following.

$$\forall X0.(v6_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v7_ordinal1\ X1)) \quad (25)$$

Assume the following.

$$\forall X0.(v1_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v1_xcmplx_0\ X1)) \quad (26)$$

Theorem 1

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(v7_ordinal1\ X1)\Rightarrow((X0 \in k2_finseq_1\ X1)\Rightarrow(k2_nat_1\ (k7_nat_d\ X1\ X0)\ np_1 \in k2_finseq_1\ X1)))$$