

t6\_nfcont\_1 (TM-  
RYvfa14n5mcuNJ2BCGhoq7orPMZ1kTdRh)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $v4\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $v2\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $l1\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k1\_normsp\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $l2\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $l1\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $l2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 \\ X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v5\_relat\_1 X1 X0))\Rightarrow(k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow(k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(((\neg v2\_struct\_0 X0)\wedge(l1\_struct\_0 \\ X0))\wedge((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 k5\_numbers (u1\_struct\_0 \\ X0))\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 \\ X0))))))\wedge(m1\_subset\_1 X2 k5\_numbers)))\Rightarrow(k1\_normsp\_1 X0 X1 X2 = \\ k1\_funct\_1 X1 X2) \end{aligned} \quad (7)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (8)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge(l1\_struct\_0 X0))\Rightarrow(\neg v1\_xboole\_0 (u1\_struct\_0 X0)) \quad (9)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (10)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (11)$$

Assume the following.

$$\forall X0.(l2\_normsp\_0 X0)\Rightarrow((l1\_normsp\_0 X0)\wedge(l2\_struct\_0 X0)) \quad (12)$$

Assume the following.

$$\forall X0.(l1\_normsp\_1 X0)\Rightarrow((l1\_rlvect\_1 X0)\wedge(l2\_normsp\_0 X0)) \quad (13)$$

Assume the following.

$$\forall X0.(l1\_normsp\_0 X0)\Rightarrow(l1\_struct\_0 X0) \quad (14)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & ((X1 \in k9\_xtuple\_0 X0) \Rightarrow ((X2 = k1\_funct\_1 X0 X1) \Leftrightarrow (k4\_tarSKI X1 X2 \in \\ & X0))) \wedge ((\neg X1 \in k9\_xtuple\_0 X0) \Rightarrow ((X2 = k1\_funct\_1 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \Rightarrow ((m1\_subset\_1 X1 X0) \Leftrightarrow \\ & (X1 \in X0))) \wedge ((v1\_xboole\_0 X0) \Rightarrow ((m1\_subset\_1 X1 X0) \Leftrightarrow (v1\_xboole\_0 \\ & X1))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (((X1 \neq k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 X2 X0 \\ & X1) \Leftrightarrow (X0 = k1\_relset\_1 X0 X2))) \wedge ((X1 = k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 \\ & X2 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (X1 = k10\_xtuple\_0 X0) \Leftrightarrow (\forall X2. (X2 \in \\ & X1) \Leftrightarrow (\exists X3. k4\_tarSKI X3 X2 \in X0)) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (X1 = k9\_xtuple\_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow \\ & (\exists X3. k4\_tarSKI X2 X3 \in X0)) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \end{aligned} \quad (21)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \end{aligned} \quad (22)$$

### Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge (v13\_algstr\_0 X0) \wedge (v2\_rlvect\_1 \\ & X0) \wedge (v3\_rlvect\_1 X0) \wedge (v4\_rlvect\_1 X0) \wedge (v5\_rlvect\_1 X0) \wedge \\ & ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v3\_normsp\_0 \\ & X0) \wedge ((v4\_normsp\_0 X0) \wedge ((v2\_normsp\_1 X0) \wedge (l1\_normsp\_1 X0)))))))))) \Rightarrow \\ & (\forall X1. ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers (u1\_struct\_0 \\ & X0)) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 \\ & X0)))))) \Rightarrow (\forall X2. (X2 \in k2\_relset\_1 (u1\_struct\_0 X0) X1) \Leftrightarrow ( \\ & \exists X3. (m2\_subset\_1 X3 k1\_numbers k5\_numbers) \wedge (X2 = k1\_normsp\_1 \\ & X0 X1 X3)))) \end{aligned}$$