

t6_ordinal3
(TMRsXApBpEp8vTRsyTjCCy43EpSoXeTQAmE)

October 27, 2020

Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_ordinal1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v3_ordinal1 X1) \Rightarrow ((X0 \in X1) \Rightarrow (v3_ordinal1 X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k2_ordinal1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow ((X2 \in X0) \wedge (v3_ordinal1 X2))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Theorem 1

$$\forall X0. (v3_ordinal1 X0) \Rightarrow (\forall X1. (r1_tarski X1 X0) \Rightarrow (k2_ordinal1 X1 = X1))$$