

t6\_pboole  
 (TMbuvqB59sV1RhL3uGUvSJubAZd8FaqepTc)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k1\_pboole : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. k1\_funct\_1 (k1\_pboole X1) X0 = k1\_xboole\_0 \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow \\ & ((\forall X3. (X3 \in X0) \Rightarrow (k1\_funct\_1 X1 X3 = k1\_funct\_1 X2 X3)) \Rightarrow (X1 = \\ & X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 (k1\_pboole X0)) \wedge ((v4\_relat\_1 (k1\_pboole X0) X0) \wedge ((v1\_funct\_1 (k1\_pboole X0)) \wedge (v1\_partfun1 (k1\_pboole X0) X0))) \quad (3)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow ((\forall X2. (X2 \in X0) \Rightarrow \\ & (k1\_funct\_1 X1 X2 = k1\_xboole\_0)) \Rightarrow (X1 = k1\_pboole X0)) \end{aligned}$$