

t6_qc_lang2

(TMMbSbL9KfwNkaZdJCqi4ERod6V7478zATR)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $v5_qc_lang1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k15_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k21_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k22_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_qc_lang1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((v5_qc_lang1 X1 X0) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 \\ X0)) \Rightarrow ((X2 = k22_qc_lang1 X0 X1) \Leftrightarrow (\exists X3.(m2_subset_1 X3 (k2_qc_lang1 \\ X0) (k3_qc_lang1 X0)) \wedge (X1 = k15_qc_lang1 X0 X3 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((v5_qc_lang1 X1 X0) \Rightarrow (\forall X2.(m2_subset_1 X2 (k2_qc_lang1 \\ X0) (k3_qc_lang1 X0)) \Rightarrow ((X2 = k21_qc_lang1 X0 X1) \Leftrightarrow (\exists X3.(\\ m1_subset_1 X3 (k9_qc_lang1 X0)) \wedge (X1 = k15_qc_lang1 X0 X2 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((v5_qc_lang1 X1 X0) \Leftrightarrow (\exists X2.(m2_subset_1 X2 (k2_qc_lang1 \\ X0) (k3_qc_lang1 X0)) \wedge (\exists X3.(m1_subset_1 X3 (k9_qc_lang1 \\ X0)) \wedge (X1 = k15_qc_lang1 X0 X2 X3)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((v5_qc_lang1 X1 X0) \Rightarrow (X1 = k15_qc_lang1 X0 (k21_qc_lang1 \\ X0 X1) (k22_qc_lang1 X0 X1)))) \end{aligned}$$