

t6_radix_2

(TMXTHc31576YWKJYtjFotyQfYYPxb8fugc5X)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_radix_1 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xreal_0 X0 k6_numbers) \wedge (r1_xreal_0 (k3_power X0 X1) k6_numbers))) \quad (2)$$

Assume the following.

$$((v2_xreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (3)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (4)$$

Assume the following.

$$\neg r1_xreal_0 np_2 np_0 \quad (5)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (6)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k1_radix_1 X0 = k3_power np_2 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (9)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg r1_xreal_0 (k1_radix_1 X0) k6_numbers)$$