

t6_real_ns1

(TMSn3QpCVFWzovroSrsZXFtDQ1XSuBGcjNH)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_card_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge ((v3_card_1 X1 X0) \wedge (v1_finseq_1 X1)))) \Rightarrow (k4_finseq_1 X1 = k2_finseq_1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow ((m2_finseq_2 X0 k1_numbers (k1_euclid (k3_finseq_1 X0))) \wedge (m1_subset_1 X0 (u1_struct_0 (k15_euclid (k3_finseq_1 X0)))))) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((k2_finseq_1 X0 = k2_finseq_1 X1) \Rightarrow (X0 = X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v1_finset_1 X0) \Rightarrow ((v1_finset_1 (k1_card_1 X0)) \wedge (v1_card_1 (k1_card_1 X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1) \wedge (v1_finseq_1 X1)) \quad (8)$$

Assume the following.

$$\forall X0.v1_card_1 (k1_card_1 X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1 X1 X0) \Leftrightarrow (k1_card_1 X1 = X0) \quad (10)$$

Assume the following.

$$\forall X0.((v3_ordinal1 X0) \wedge (v1_finset_1 X0)) \Rightarrow (v7_ordinal1 X0) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0))) \quad (12)$$

Assume the following.

$$\forall X0.(v1_card_1 X0) \Rightarrow (v3_ordinal1 X0) \quad (13)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m2_finseq_1 X1 k1_numbers) \Rightarrow ((k4_finseq_1 X1 = k2_finseq_1 X0) \Rightarrow (m2_finseq_2 X1 k1_numbers (k1_euclid X0))))$$