

t6_rfinseq
(TMKaSrEMgSdcsTZmqHdLqqijUN8N27tMVz)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k17_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k16_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski X0 (k9_xtuple_0 X1)) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = X0)) \quad (1)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_tarski (k2_finseq_1 X0) (k2_finseq_1 X1)))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k5_relat_1 X2 X1)) \Rightarrow (k1_funct_1 (k5_relat_1 X2 X1) X0 = k1_funct_1 X2 X0)) \quad (3)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X0 \in k2_finseq_1 X1) \Leftrightarrow ((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 X1)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v7_ordinal1 X1) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k17_finseq_1 X0 X1 X2 = k16_finseq_1 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (10)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow ((v1_finseq_1 X0) \Leftrightarrow (\exists X1. (v7_ordinal1 X1) \wedge (k9_xtuple_0 X0 = k2_finseq_1 X1))) \quad (11)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow (k16_finseq_1 X0 X1 = k5_relat_1 X1 (k2_finseq_1 X0))) \quad (12)$$

Theorem 1

$$\forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m2_finseq_1 X1 X0) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 X3) \Rightarrow ((X2 \in k4_finseq_1 X1) \wedge (X3 \in k2_finseq_1 X2)) \Rightarrow ((k1_funct_1 (k17_finseq_1 X0 X2 X1) X3 = k1_funct_1 X1 X3) \wedge (X3 \in k4_finseq_1 X1))))))$$