

t6_rfunct_1
(TMHN3aytDUPonobVmuz5c7yzMHwiHzYD3et)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_rfunct_1 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow (k8_relat_1 (k4_rfunct_1 X0) (k1_tarski k6_numbers) = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. k4_xboole_0 X0 k1_xboole_0 = X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (6)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (k5_relat_1 X0 (k9_xtuple_0 X0) = X0) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ ((v1_relat_1 (k4_rfunct_1 X0)) \wedge ((v1_funct_1 (k4_rfunct_1 X0)) \wedge \\ (v1_valued_0 (k4_rfunct_1 X0)))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.m1_subset_1 (k6_subset_1 X0 X1) (k1_zfmisc_1 X0) \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ ((v1_relat_1 (k4_rfunct_1 X0)) \wedge (v1_funct_1 (k4_rfunct_1 X0))) \end{aligned} \quad (10)$$

Assume the following.

$$k1_xboole_0 = the (\lambda X0 : \iota.v1_xboole_0 X0) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X1 = k4_rfunct_1 \\ X0) \Leftrightarrow ((k9_xtuple_0 X1 = k6_subset_1 (k9_xtuple_0 X0) (k8_relat_1 \\ X0 (k1_tarski k6_numbers)) \wedge (\forall X2.(X2 \in k9_xtuple_0 X1) \Rightarrow \\ (k1_funct_1 X1 X2 = k5_xcmplx_0 (k1_funct_1 X0 X2)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (13)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ (k9_xtuple_0 (k4_rfunct_1 (k4_rfunct_1 X0)) = k9_xtuple_0 (k5_relat_1 \\ X0 (k9_xtuple_0 (k4_rfunct_1 X0)))) \end{aligned}$$