

## t6\_sheffer2

(TMR9WddTEAjGoA2RtpQrKwA8fybjumESfEq)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v1\_sheffer2 : \iota \Rightarrow o$  be given. Let  $l1\_sheffer1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v1\_sheffer2 X0) \wedge (l1\_sheffer1 \\ X0))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (k5\_sheffer1 \\ X0 X1 (k5\_sheffer1 X0 (k5\_sheffer1 X0 X1 X1) X1) = k5\_sheffer1 X0 X1 \\ X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (((\neg v2\_struct\_0 X0) \wedge (l1\_sheffer1 \\ X0)) \wedge ((m1\_subset\_1 X1 (u1\_struct\_0 X0)) \wedge (m1\_subset\_1 X2 (u1\_struct\_0 \\ X0)))) \Rightarrow (m1\_subset\_1 (k5\_sheffer1 X0 X1 X2) (u1\_struct\_0 X0)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge (l1\_sheffer1 X0)) \Rightarrow ((v1\_sheffer2 \\ X0) \Leftrightarrow (\forall X1. (m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. \\ (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3. (m1\_subset\_1 X3 \\ (u1\_struct\_0 X0)) \Rightarrow (k5\_sheffer1 X0 (k5\_sheffer1 X0 X1 (k5\_sheffer1 \\ X0 (k5\_sheffer1 X0 X2 X1) X1)) (k5\_sheffer1 X0 X2 (k5\_sheffer1 X0 \\ X3 X1)) = X2)))))) \end{aligned} \tag{3}$$

### Theorem 1

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v1\_sheffer2 X0) \wedge (l1\_sheffer1 \\ X0))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (k5\_sheffer1 \\ X0 (k5\_sheffer1 X0 X1 (k5\_sheffer1 X0 (k5\_sheffer1 X0 X1 X1) X1)) \\ (k5\_sheffer1 X0 X1 X1) = X1)) \end{aligned}$$