

t6_sin_cos7

(TMMp9HE5A6vp4xgecY7oPcg2czmiWJ7NBNy)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k12_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 \\ & X1 X2)) \Rightarrow (r1_xxreal_0 X1 (k2_xcmplx_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 \\ & k6_numbers X0) \wedge (r1_xxreal_0 k6_numbers X1)) \Rightarrow (r1_xxreal_0 k6_numbers \\ & (k7_xcmplx_0 X0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (4)$$

Assume the following.

$$k2_xcmplx_0 np_1 (k4_xcmplx_0 np_1) = np_0 \quad (5)$$

Assume the following.

$$r1_xxreal_0 np_0 np_1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k9_binop_2 X0 X1 = k2_xcmplx_0 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k12_binop_2 X0 X1 = k7_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(\forall X2. (v1_xreal_0 X2)\Rightarrow(((r1_xxreal_0 X0 X1)\wedge(r1_xxreal_0 X1 X2))\Rightarrow(r1_xxreal_0 X0 X2)))) \quad (9)$$

Assume the following.

$$k2_xcmplx_0 np_1 (k4_xcmplx_0 np_1) = k6_numbers \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (12)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(((r1_xxreal_0 k6_numbers X0)\wedge(r1_xxreal_0 np_1 X1))\Rightarrow(r1_xxreal_0 k6_numbers (k12_binop_2 (k9_binop_2 X1 np_1) X0))))$$