

t6_tex_4 (TMMrDd-
kwFRqaZ3sdusgpzUB1wV56nhxMJuP)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_tex_4 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v3_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (2)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \quad (3)$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (l1_struct_0 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m1_subset_1 (k6_domain_1 X0 X1) (k1_zfmisc_1 X0)) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow ((v1_tex_4 X1 X0) \Leftrightarrow (\forall X2. (m1_subset_1 \\ X2 (u1_struct_0 X0)) \Rightarrow (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow (((v3_pre_topc X3 X0) \wedge ((X2 \in X3) \wedge (X2 \in X1))) \Rightarrow \\ (r1_tarski X1 X3)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarSKI X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (7)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (v1_tex_4 (k6_domain_1 (u1_struct_0 X0) X1) X0))$$