

t6_tops_2 (TMHf- sWqfXxdmF3j7XpwaUH4KHtP7jUEpFck)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k6_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_subset_1 : \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((X1 \neq k1_xboole_0) \Rightarrow (k7_subset_1 X0 (k2_subset_1 X0) (k5_setfam_1 X0 X1) = k6_setfam_1 X0 (k7_setfam_1 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (m1_subset_1 (k5_setfam_1 X0 X1) (k1_zfmisc_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0. m1_subset_1 (k2_subset_1 X0) (k1_zfmisc_1 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 X1 = k4_xboole_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0. k2_subset_1 X0 = X0 \quad (6)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((X1 \neq k1_xboole_0) \Rightarrow (k6_setfam_1 X0 (k7_setfam_1 X0 X1) = k3_subset_1 X0 (k5_setfam_1 X0 X1)))$$