

t6_xxreal_2
(TMR8JttdkvfiTsRf7tVf4siSLnYhFg16bjZ)

October 27, 2020

Let $v2_membered : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_xxreal_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(m1_xxreal_2 X1 X0) \Rightarrow (v1_xxreal_0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((m1_xxreal_2 X1 X0) \Leftrightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((X2 \in X0) \Rightarrow (r1_xxreal_0 X2 X1))))) \quad (3)$$

Theorem 1

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v2_membered X1) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (\forall X2.(m1_xxreal_2 X2 X1) \Rightarrow (m1_xxreal_2 X2 X0))))$$