

t6_yellow_6
(TMVnNg8s9zMEAy99rJxQVbwz2F3Qu3ijhUV)

October 27, 2020

Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $v4_yellow_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_yellow_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k1_toler_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_wellord1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_orders_2 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (k1_toler_1 X0 X1 = k2_wellord1 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. (l1_orders_2 X0) \Rightarrow (m1_subset_1 (u1_orders_2 X0) (k1_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X0)))) \quad (5)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. k2_wellord1 X0 X1 = k3_xboole_0 X0 (k2_zfmisc_1 X1 X1)) \quad (6)$$

Assume the following.

$$\forall X0. (l1_orders_2 X0) \Rightarrow (\forall X1. (m1_yellow_0 X1 X0) \Rightarrow ((v4_yellow_0 X1 X0) \Leftrightarrow (u1_orders_2 X1 = k1_toler_1 (u1_orders_2 X0) (u1_struct_0 X1)))) \quad (7)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0) \Rightarrow (\forall X1.(l1_orders_2 X1) \Rightarrow ((m1_yellow_0 X1 X0) \Leftrightarrow ((r1_tarski (u1_struct_0 X1) (u1_struct_0 X0)) \wedge (r1_tarski (u1_orders_2 X1) (u1_orders_2 X0)))))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (9)$$

Theorem 1 $\forall X0.(l1_orders_2 X0) \Rightarrow ((v4_yellow_0 X0 X0) \wedge (m1_yellow_0 X0 X0)).$