

t70_eqrel_1

(TMJCx2n8D1RCT6sxQwtC3YGwhrreSFVw4kn)

October 27, 2020

Let $m1_eqrel_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2.\neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2.(X2 \in X0) \wedge (X2 \in X1)) \wedge (r1_xboole_0 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m1_eqrel_1 X1 X0) \Rightarrow (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((m1_eqrel_1 X1 X0) \Leftrightarrow ((k5_setfam_1 X0 X1 = X0) \wedge (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((X2 \in X1) \Rightarrow ((X2 \neq k1_xboole_0) \wedge (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 X0)) \Rightarrow (\neg(X3 \in X1) \wedge ((X2 \neq X3) \wedge (\neg r1_xboole_0 X2 X3)))))))))) \quad (4)$$

Theorem 1

$$\forall X0.\forall X1.(m1_eqrel_1 X1 X0) \Rightarrow (\forall X2.\forall X3.\forall X4.((X2 \in X3) \wedge ((X3 \in X1) \wedge ((X2 \in X4) \wedge (X4 \in X1)))) \Rightarrow (X3 = X4))$$