

t70_xboolean
(TMae2SnrTbhvp2Wcuex2QQJRFcF8eusM2US)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k9_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k9_xboolean X0 (k7_xboolean X0 X1) = k4_xboolean (k3_xboolean X0) X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (v1_xboolean (k3_xboolean X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k11_xboolean X0 X1 = k4_xboolean X0 (k3_xboolean X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (k4_xboolean X0 X1 = k4_xboolean X1 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k9_xboolean X0 (k7_xboolean X0 X1) = k11_xboolean X1 X0))$$