

t71\_finseq\_2  
(TMbvDdvBDzzn7cQKajx3j3UEGq5iPq1EJHL)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((r1\_tarski X0 X1) \wedge (r1\_tarski X2 X3)) \Rightarrow (r1\_tarski (k2\_zfmisc\_1 X0 X2) (k2\_zfmisc\_1 X1 X3)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2))) \Rightarrow (\forall X3. ((v1\_relat\_1 X3) \wedge (v1\_funct\_1 X3)) \Rightarrow (((r1\_tarski \\ & (k2\_zfmisc\_1 (k10\_xtuple\_0 X0) (k10\_xtuple\_0 X1)) (k9\_xtuple\_0 X3)) \wedge (X2 = k3\_funcop\_1 X3 X0 X1)) \Rightarrow (k3\_finseq\_1 X2 = k3\_xxreal\_0 \\ & (k3\_finseq\_1 X0) (k3\_finseq\_1 X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge(v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1)))\Rightarrow((m1\_finseq\_1 X1 X0)\Leftrightarrow(r1\_tarski (k10\_xtuple\_0 X1) X0)) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 X1)))\Rightarrow(((X1\neq k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 \\ X1)\Leftrightarrow(X0 = k1\_relset\_1 X0 X2)))\wedge((X1 = k1\_xboole\_0)\Rightarrow((v1\_funct\_2 \\ X2 X0 X1)\Leftrightarrow(X2 = k1\_xboole\_0)))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (10)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(\neg v1\_xboole\_0 X1)\Rightarrow \\ (\forall X2.(\neg v1\_xboole\_0 X2)\Rightarrow(\forall X3.((v1\_relat\_1 X3)\wedge \\ ((v1\_funct\_1 X3)\wedge(v1\_finseq\_1 X3)))\Rightarrow(\forall X4.((v1\_funct\_1 \\ X4)\wedge((v1\_funct\_2 X4 (k2\_zfmisc\_1 X0 X1) X2)\wedge(m1\_subset\_1 X4 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1) X2))))\Rightarrow(\forall X5.(m2\_finseq\_1 \\ X5 X0)\Rightarrow(\forall X6.(m2\_finseq\_1 X6 X1)\Rightarrow((X3 = k3\_funcop\_1 X4 X5 \\ X6)\Rightarrow(k3\_finseq\_1 X3 = k3\_xreal\_0 (k3\_finseq\_1 X5) (k3\_finseq\_1 \\ X6)))))))))) \end{aligned}$$