

t71_funct_1
(TMSD9pXaa6HpbhoGx9TCV4C2ksKkxQbfyxy)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1)) \wedge (r1_xboole_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((k4_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (X1 = k1_funct_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X0 \in k8_relat_1 X2 X1) \Leftrightarrow (\exists X3. (X3 \in k10_xtuple_0 X2) \wedge ((k4_tarski X0 X3 \in X2) \wedge (X3 \in X1)))) \quad (3)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. (r1_xboole_0 X1 X2) \Rightarrow (r1_xboole_0 (k8_relat_1 X0 X1) (k8_relat_1 X0 X2)))$$