

t71_qc_lang3

(TMPz8q1WfgBpD1HckwoQL2RwApQMNnXS7Ns)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k6_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k4_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\
 & X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0)) \Rightarrow (\forall X3. \\
 & ((\neg v1_xboole_0 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_qc_lang1 \\
 & X0)))) \Rightarrow (k4_qc_lang3 X0 X3 (k4_qc_lang2 X0 X1 X2) = k4_subset_1 X3 \\
 & (k4_qc_lang3 X0 X3 X1) (k4_qc_lang3 X0 X3 X2)))))) \\
 & \tag{1}
 \end{aligned}$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\neg v1_xboole_0 (k4_qc_lang1 X0)) \tag{2}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0) \wedge ((m1_subset_1 \\
 & X1 (k9_qc_lang1 X0)) \wedge (m1_subset_1 X2 (k9_qc_lang1 X0)))) \Rightarrow (m1_subset_1 \\
 & (k4_qc_lang2 X0 X1 X2) (k9_qc_lang1 X0)) \\
 & \tag{3}
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k4_qc_lang1 X0) (k1_zfmisc_1 \\
 & (k2_qc_lang1 X0))) \\
 & \tag{4}
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\
 & X0)) \Rightarrow (k6_qc_lang3 X0 X1 = k4_qc_lang3 X0 (k4_qc_lang1 X0) X1)) \\
 & \tag{5}
 \end{aligned}$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1\ X0)) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k9_qc_lang1\ X0)) \Rightarrow (k6_qc_lang3 \\ X0\ (k4_qc_lang2\ X0\ X1\ X2) = k4_subset_1\ (k4_qc_lang1\ X0)\ (k6_qc_lang3\\ X0\ X1)\ (k6_qc_lang3\ X0\ X2)))) \end{aligned}$$