

t71_relat_1 (TMKJ-
DuneSxKA23XA3jNRLuMkSSGBLioAxaG)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarSKI : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (v1_relat_1 (k5_relat_1 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \quad (2)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. (v1_relat_1 X1) \Rightarrow ((X0 = X1) \Leftrightarrow (\forall X2. \forall X3. (k4_tarSKI X2 X3 \in X0) \Leftrightarrow (k4_tarSKI X2 X3 \in X1)))) \quad (3)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X2 = k5_relat_1 X0 X1) \Leftrightarrow (\forall X3. \forall X4. (k4_tarSKI X3 X4 \in X2) \Leftrightarrow ((X3 \in X1) \wedge (k4_tarSKI X3 X4 \in X0))))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow (k5_relat_1 (k5_relat_1 X2 X0) X1 = k5_relat_1 X2 (k3_xboole_0 X0 X1))$$