

## t71\_rinfsup1

(TMRn4sBvKtG5oYtmxUCfnRb2zfvkU58XQbd)

October 27, 2020

Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v8\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v2\_seq\_2 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_seq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_rinfsup1 : \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_seq\_2 : \iota \Rightarrow o$  be given. Let  $k1\_rinfsup1 : \iota \Rightarrow \iota$  be given. Let  $v1\_comseq\_2 : \iota \Rightarrow o$  be given. Let  $k4\_rinfsup1 : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xbool\_0 : \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v3\_membered : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 \\ & X1 k5\_numbers k1\_numbers) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & k5\_numbers k1\_numbers)))))) \Rightarrow ((\forall X2.(m2\_subset\_1 X2 k1\_numbers \\ & k5\_numbers) \Rightarrow (r1\_xxreal\_0 (k1\_seq\_1 X1 X2) X0)) \Leftrightarrow ((v1\_seq\_2 X1) \wedge \\ & (r1\_xxreal\_0 (k1\_rinfsup1 X1) X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow \\ & ((v1\_comseq\_2 X0) \Rightarrow ((v1\_comseq\_2 (k4\_rinfsup1 X0)) \wedge (v1\_comseq\_2 \\ & (k3\_rinfsup1 X0)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\ & ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers k1\_numbers) \wedge (m1\_subset\_1 \\ & X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow ((v1\_comseq\_2 \\ & X1) \Rightarrow (r1\_xxreal\_0 (k1\_rinfsup1 (k3\_rinfsup1 X1)) (k1\_seq\_1 (k4\_rinfsup1 \\ & X1) X0)))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow \\ & (((v8\_valued\_0 X0) \wedge (v2\_seq\_2 X0)) \Rightarrow (v1\_comseq\_2 X0)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2\_xreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v3\_valued\_0 \\ & X0))) \Rightarrow (k1\_seq\_1 X0 X1 = k1\_funct\_1 X0 X1) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. (m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\ & ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers k1\_numbers) \wedge (m1\_subset\_1 \\ & X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow ((v8\_valued\_0 \\ & X1) \Rightarrow (k1\_seq\_1 (k4\_rinf sup1 X1) X0 = k1\_seq\_1 X1 X0)) \end{aligned} \quad (9)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v3\_valued\_0 \\ & X0))) \Rightarrow (v1\_xreal\_0 (k1\_funct\_1 X0 X1)) \end{aligned} \quad (11)$$

Assume the following.

$$v3\_membered k1\_numbers \quad (12)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (13)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow \\ & ((v1\_funct\_1 (k3\_rinf sup1 X0)) \wedge ((v1\_funct\_2 (k3\_rinf sup1 X0) \\ & k5\_numbers k1\_numbers) \wedge (m1\_subset\_1 (k3\_rinf sup1 X0) (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 X1)) \Rightarrow (m2\_subset\_1 (k2\_nat\_1 X0 X1) k1\_numbers k5\_numbers) \quad (16)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 X1)) \Rightarrow (k2\_nat\_1 X0 X1 = k2\_nat\_1 X1 X0) \quad (17)$$

Assume the following.

$$\forall X0. (m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v3\_valued\_0 \\ & X0) \wedge (v1\_comseq\_2 X0)))) \Rightarrow ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge \\ & ((v3\_valued\_0 X0) \wedge ((v1\_seq\_2 X0) \wedge (v2\_seq\_2 X0)))))) \end{aligned} \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (20)$$

Assume the following.

$$\forall X0. \forall X1. (v3\_membered X1) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v3\_valued\_0 X2)) \quad (21)$$

### Theorem 1

$$\begin{aligned} & \forall X0. (m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\ & ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers k1\_numbers) \wedge (m1\_subset\_1 \\ & X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow ((( \\ & v8\_valued\_0 X1) \wedge (v2\_seq\_2 X1)) \Rightarrow (r1\_xxreal\_0 (k1\_seq\_1 (k3\_rinf sup1 \\ & X1) (k2\_nat\_1 X0 np\_1)) (k1\_seq\_1 X1 X0)))) \end{aligned}$$