

t71_xreal_1
(TMdSZLSNCa1Qiv1yXB1twxqbK3eexQbS3Qd)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & k6_numbers X0) \wedge ((r1_xxreal_0 X0 X1) \wedge ((r1_xxreal_0 k6_numbers \\ & X2) \wedge (r1_xxreal_0 X2 X3)))) \Rightarrow (r1_xxreal_0 (k3_xcmplx_0 X0 X2) (\\ & k3_xcmplx_0 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \Rightarrow (r1_xxreal_0 X1 (k2_xcmplx_0 X0 X1)))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 k6_numbers = k6_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 k6_numbers = X0) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg \\ & r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 X1 k6_numbers) \wedge (r1_xxreal_0 \\ & (k3_xcmplx_0 X0 X1) k6_numbers)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k3_xcmplx_0 X0 X1)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (\neg(r1_xxreal_0 \\ & k6_numbers X0) \wedge ((r1_xxreal_0 k6_numbers X1) \wedge ((r1_xxreal_0 k6_numbers \\ & X2) \wedge ((r1_xxreal_0 k6_numbers X3) \wedge ((k2_xcmplx_0 (k3_xcmplx_0 \\ & X0 X2) (k3_xcmplx_0 X1 X3) = k6_numbers) \wedge ((X0 \neq k6_numbers) \wedge (X2 \neq \\ & k6_numbers)))))))))) \end{aligned}$$