

t71_xxreal_1
(TMLk8mB4xKtK3yM1Gi5jaqgaVfAzB1bba5X)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_tarski (k1_xxreal_1 X0 X1) (k2_xxreal_1 X2 X3))) \Rightarrow ((\\ & r1_xxreal_0 X2 X0) \wedge (\neg r1_xxreal_0 X3 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_tarski (\\ & k2_xxreal_1 X0 X1) (k1_xxreal_1 X2 X3)) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (\\ & r1_xxreal_0 X2 X0) \wedge (r1_xxreal_0 X1 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (\neg (r1_xxreal_0 \\ & X0 X1) \wedge (k1_xxreal_1 X0 X1 = k2_xxreal_1 X2 X3)))))) \end{aligned}$$