

t73_mcart_1

(TMKPCz2rP9X1CEZ6Qe5sgywH9fEf9PtptmN)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((r1_tarski X0 X1) \wedge (r1_tarski X2 X3)) \Rightarrow (r1_tarski (k2_zfmisc_1 X0 X2) (k2_zfmisc_1 X1 X3)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. k3_zfmisc_1 X0 X1 X2 = k2_zfmisc_1 (k2_zfmisc_1 X0 X1) X2 \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. ((r1_tarski X0 X1) \wedge ((r1_tarski X2 X3) \wedge (r1_tarski X4 X5))) \Rightarrow (r1_tarski (k3_zfmisc_1 X0 X2 X4) (k3_zfmisc_1 X1 X3 X5))$$